

Gravitational Waves from Spinning Non-Abelian Cosmic Strings

R. J. Slagter

*University of Amsterdam, Physics Department, The Netherlands,
SARA, Stichting Academisch Rekencentrum Amsterdam,
and
ASFYON, Astronomisch Fysisch Onderzoek Nederland**

We investigated the SU(2) Einstein-Yang-Mills system on a time-dependent non-diagonal cylindrical symmetric spacetime. From the numerical investigation, wave-like solutions are found, consistent with the familiar string-like features. They possess an angle deficit which depends on the initial form of the magnetic component of the Yang-Mills field, i.e., the number of times it crosses the r-axis. The appearance of a curvature singularity depends on the ratio of the Planck scale M_{pl} and Yang-Mills couplings constant g , i.e., $\alpha = M_{pl}/g = \frac{1}{\sqrt{4\pi G}g}$ and the initial mass per unit length of the cosmic string. The solitonlike behavior of gravitational and Yang-Mills waves show significantly differences from the ones found in the Abelian Einstein-Maxwell system. The stability of the resulting non-Abelian spinning string is analyzed using the multiple-scale method. To first order a consistent set of equations is obtained.

However, a throughout analysis is necessary in order to conclude if the solution is topological stable and will have magnetic charge.

I. INTRODUCTION

Taking the Yang-Mills model as the matter part of the equations of Einstein, one finds a rich spectrum of stationary regular and non-regular solutions. This in contrast with the vacuum solution and Einstein-Maxwell(EM) solutions, where the Schwarzschild respectively the Reissner-Nordström(RN) solution are the only static black hole solutions. In the Einstein-Yang-Mills(EYM) theory one finds not only the embedding of the Abelian RN black hole, but in addition there are genuine non-Abelian coloured black holes. Their co-existence gives rise to the violation of the no-hair conjecture, since they carry the same (magnetic) charge. For an overview, see Volkov and Gal'tsov [1]. After the discovery by Bartnik and McKinnon(BMK) [2] of these non-Abelian particle-like and later the non-RN (hairy) black hole solutions of the SU(2) EYM theory [3], throughout instability investigations where done by several authors [4–8]. It turns out that under spherically symmetric perturbations the BMK solution is unstable. The n th BMK solution has in fact $2n$ unstable modes, comparable with the flat space-time 'sphaleron' solution. In the static case, it was recently conjectured [9] that there are hairy black holes in the SU(N) EYM-theories with topological instabilities. In order to analyze the stability of the solutions, one usually linearized the field equations [4,6]. We conjectured that the conventional linear analysis is in fact inadequate to be applied to the situation where a singularity is formed [10]. One better can apply the so-called multiple-scale or two-timing method, developed by Choquet-Bruhat [11–13]. This method is particularly useful for constructing uniformly valid approximations to solutions of perturbation problems [14]. At the threshold of black hole formation it is found numerically [15–18] that there is a critical parameter whose value separates solutions containing black holes from those which do not. For a critical value one observes self-similarity: it produces itself (echoes) on progressively finer scales (Choquet-Bruhat-scaling). It is evident that the collapsing ball of field energy will produce gravitational waves, which will be coupled to the YM-field perturbations. Due to the accumulation of echoes, the curvature diverges. Close related to this critical behavior is the comparable irregularities which can be found in the Einstein-Skyrme(ES) model and the Einstein-Yang-Mills-Higgs(EYMH) model. In the latter model, space-like hypersurfaces develop for a critical value of $\alpha = M_W/gM_{pl}$ two distinct regions separated by a long throat, with M_W the YM scale and $M_{pl} = 1/\sqrt{4\pi G}$.

The solutions mentioned above need not be spherically symmetric. Since the pioneering work of Stachel [19], we know that stationary cylindrical symmetric rotating models possess intriguing features. First, the evolution of the 'classical' cylindrical gravitational wave-solution in the vacuum situation can be described in terms of two effects: the usual cylindrical reflections and a rotation of the polarization vector of the waves, an effect comparable with the Faraday rotation [20]. Secondly, the time-dependent solutions can be related to the cosmic-string solution interacting with gravitational waves [21]. Usually the cosmic string solution, with its famous conical structure, is found in the

*Electronic address: rjs@asfyon.nl

U(1)-gauge Einstein-Higgs system. It is remarkable that the cylindrical rotating vacuum solution possesses the same conical structure. Moreover, it is asymptotically flat and free of singularities. The physical interpretation can nicely be formulated using the Belinsky-Zakharov method of integrating the Einstein equations with the help of the Ernst potentials [22]. Using the C-energy description, one can conclude that the solution represents solitonic gravitational waves interacting with a straight-line cosmic string, where the asymptotic angle deficit depends on the C-energy of the gravitational waves. When a Maxwell field is incorporated, the physical interpretation of the solutions is not easily formulated. It is hard to believe that in the electrovac solution the electro-magnetic waves do not contribute to the angle deficit [23]. Moreover, the mass per unit length of the cosmic string, which is a constant in the Xanthopoulos solution, will be determined by the energy-momentum tensor of the non-vacuum situation. In stead of 'electrifying' or 'magnetifying' the vacuum solutions via the Ernst formulation, one can better consider the coupled system of Einstein equations and matter field equations.

Here we consider the Einstein-Yang-Mills system on a cylindrical symmetric rotating space-time. This situation can formally be obtained from the axially symmetric situation by complex substitution of $t \rightarrow iz, z \rightarrow it$ [24]. In the non-rotating cylindrically symmetric situation, it was numerically found [26] that the singular behavior at finite distance of the core as found in the supermassive situation, will be pushed to infinity for some critical values of the gauge coupling constant and one of the YM components.

We will use a (2+1)+1 reduction scheme [25] in order to obtain the suitable parameterization for the Yang-Mills potential. In fact one replaces the YM field by a 2D Abelian one-form, a 2D Kaluza-Klein two-form, a 2D matrix-valued scalar and a 3D Higgs field [27].

In the Abelian counterpart model it is known that the space-time around (spinning) gauge strings could have, besides the usual angle deficit feature, exotic properties, such as the violation of causality [28,29], time-delay effects [30], helical structure of time [31–33] and frame-dragging [34]. A class of approximate solutions of the coupled Einstein-scalar-gauge field equations on the (r, z) -plane of an axially symmetric space-time was found [35]. So a question addressed in this paper will be whether the string features will be maintained in the EYM system. In the general (r, z) -dependent situation, a nice overview was given by Islam [36]. One can proof via the Ernst formulation that for any solution of the stationary Einstein equations one can generate a corresponding solution of the EM equations. In fact, one introduces two complex potentials, for which one finds two second-order elliptic differential equations. Starting with the Kerr solution, it is possible to use this correspondence to find a solution of the EM equations leading to the Kerr-Newman solution. This 'Ernst-route' fails in the EYM system, simply by the fact that one of the metric components does not decouple from the other ones. It is also known that the electrovac staticity theorem [37] does not generalize to the EYM system. In fact, spinning EYM systems must be electrically charged. So another question addressed in this paper is whether electrically and magnetically charged solution will exist in the EYM system. Finally, this is not the complete story: the investigations can be extended to the model where the interior solution is properly matched onto the exterior solution of the string [38]. This is currently under study by the author.

The plan of this paper is as follows. In Sec. II we derive the field equation on an cylindrical symmetric space time. In Sec. III we present the numerical solution. In section IV we present a approximate solution using the multiple-scale method and in Sec. V we summarize and analyse our results.

II. THE FIELD EQUATIONS OF THE EINSTEIN YANG-MILLS SYSTEM

Consider the Lagrangian of the SU(2) EYM system

$$S = \int d^4x \sqrt{-g} \left[\frac{\mathcal{R}}{16\pi G} - \frac{1}{4} \mathcal{F}_{\mu\nu}^a \mathcal{F}^{\mu\nu a} \right], \quad (1)$$

with the YM field strength

$$\mathcal{F}_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g\epsilon^{abc} A_\mu^b A_\nu^c, \quad (2)$$

g the gauge coupling constant, G Newton's constant, A_μ^a the gauge potential, and $\mathcal{R}$ the curvature scalar. The field equations then become

$$G_{\mu\nu} = 8\pi G \mathcal{T}_{\mu\nu}, \quad (3)$$

$$\mathcal{D}_\mu \mathcal{F}^{\mu\nu a} = 0, \quad (4)$$

with $\mathcal{T}$ the energy-momentum tensor

$$\mathcal{T}_{\mu\nu} = \mathcal{F}_{\mu\lambda}^a \mathcal{F}_\nu^{\lambda a} - \frac{1}{4} g_{\mu\nu} \mathcal{F}_{\alpha\beta}^a \mathcal{F}^{\alpha\beta a}, \quad (5)$$

and $\mathcal{D}$ the gauge-covariant derivative,

$$\mathcal{D}_\alpha \mathcal{F}_{\mu\nu}^a \equiv \nabla_\alpha \mathcal{F}_{\mu\nu}^a + g \epsilon^{abc} A_\alpha^b \mathcal{F}_{\mu\nu}^c. \quad (6)$$

Let us consider the cylindrical space time

$$ds^2 = -\Omega^2(dt^2 - dr^2) + f(dz + \omega d\varphi)^2 + \frac{r^2}{f} d\varphi^2, \quad (7)$$

where Ω, f and ω are functions of t and r . The problem is, which ansatz one needs for the YM field. It is often argued that Einsteins gravity on a three dimensional space time shows some similarity with Chern Simons theory with the Poincaré group being the underlying gauge group. So when trying to find the dimensional reduction for the stationary axially symmetric EYM system, one can be guided by the results found by using the (2+1)+1 reduction scheme suggested by Maeda, Sasaki, Nakamura and Miyama [25], and later worked out by Gal'tsov [27]. One writes in general

$$ds^2 = -e^\psi(dt + v_i dx^i)^2 + e^{2\phi-\psi}(d\varphi + k_a dx^a)^2 + g_{ab} dx^a dx^b, \quad (8)$$

where $v_i dx^i = \omega(d\varphi + k_a dx^a) + \nu_a dx^a$, k_a an one-form, generating a field strength $\kappa_{ab} = \partial_a k_b - \partial_b k_a$. The one-form ν_a give rise to the 2D field strength $\omega_{ab} = \partial_a \nu_b - \partial_b \nu_a + \omega \kappa_{ab}$ ($a, b=1, 2$). The YM potential is then written as

$$A_\mu dx^\mu = \mathbf{a}_a dx^a + \mathbf{\Phi}(dt + \nu_i dx^i) + \mathbf{\Psi}(d\varphi + k_a dx^a) \quad (9)$$

Here $\mathbf{a}$ is the dynamical part of the YM field, parameterized by a 2D complex Abelian one-form, $\mathbf{\Phi}$ a 3D Higgs field and $\mathbf{\Psi}$ a matrix-valued scalar. In our simplified case of metric (7), we have $\nu_1 = -\omega k_1, k_2 = \nu_2 = 0$ and the YM potential can be as

$$A_\mu = \mathbf{\Phi}(dz + \omega d\varphi) + \mathbf{\Psi}(d\varphi - k_1 dt) + \mathbf{a}_a dx^a \quad (10)$$

The YM-part of the Lagrangian can then be written as

$$\begin{aligned} \mathcal{L}_{YM} = \Omega^2 r \Big[& \left\{ f_{ab} f^{ab} + \frac{2f}{r^2} (D_a \mathbf{\Psi} + \mathbf{\Phi} \partial_a \omega - a'_a)(D^a \mathbf{\Psi} + \mathbf{\Phi} \partial^a \omega - a'^a) \right\} \\ & + \frac{2}{f} \left\{ D_a \mathbf{\Phi} D^a \mathbf{\Phi} + \frac{f}{r^2} (\mathbf{\Phi}' + g[\mathbf{\Phi}, \mathbf{\Psi}])^2 \right\} \Big], \end{aligned} \quad (11)$$

with

$$D_a \mathbf{\Psi} = \partial_a \mathbf{\Psi} + g[a_a, \mathbf{\Psi}] - k_a \mathbf{\Psi}', \quad (12)$$

f_{ab} the 2D field strength

$$f_{ab} = \partial_a a_b - \partial_b a_a + g[a_a, a_b] + a'_a k_b - a'_b k_a + \mathbf{\Phi} \omega_{ab} + \mathbf{\Psi} k_{ab}, \quad (13)$$

and a prime denoting the partial derivative with respect to φ .

Let us consider the following parameterization

$$\begin{aligned} a_1 &= A_2(t, r) \tau_\varphi, & a_2 &= A_1(t, r) \tau_\varphi, & \mathbf{\Psi} &= W_1(t, r) \tau_r + (W_2(t, r) - 1) \tau_z, \\ \mathbf{\Phi} &= \Phi_1(t, r) \tau_r + \Phi_2(t, r) \tau_z, \end{aligned} \quad (14)$$

with τ_i the usual set of orthonormal vectors. From the condition $T_{tt} - T_{rr} = 0$, we obtain

$$\partial_t A_1 = \partial_r A_2, \quad W_2 = \frac{\Phi_2 W_1}{\Phi_1} + \frac{g-1}{g}. \quad (15)$$

From the Einstein equations and the YM equations we obtain the set of partial differential equations for $\Omega, f, \omega, \Phi_1, \Phi_2$ and W_1 . It follows from a combination of the YM equations, that A_1 and A_2 can be expressed in Φ_1 and Φ_2 :

$$A_1 = \frac{\Phi_1 \partial_r \Phi_2 - \Phi_2 \partial_r \Phi_1}{g(\Phi_1^2 + \Phi_2^2)}, \quad A_2 = \frac{\Phi_1 \partial_t \Phi_2 - \Phi_2 \partial_t \Phi_1}{g(\Phi_1^2 + \Phi_2^2)} \quad (16)$$

For the gauge $\partial_t A_2 = \partial_r A_1$ and hence $\partial_t^2 A_i - \partial_r^2 A_i = 0$, we obtain from the Einstein equations Eq.(3)

$$\partial_t^2 \Omega - \partial_r^2 \Omega + \frac{f^2 \Omega}{4r^2} [(\partial_t \omega)^2 - (\partial_r \omega)^2] + \frac{1}{\Omega} [(\partial_r \Omega)^2 - (\partial_t \Omega)^2] + \frac{\Omega}{4f^2} [(\partial_t f)^2 - (\partial_r f)^2] + \frac{\Omega}{2fr} \partial_r f = 0, \quad (17)$$

$$\begin{aligned} & \partial_t^2 f - \partial_r^2 f + \frac{1}{f} [(\partial_r f)^2 - (\partial_t f)^2] + \frac{f^3}{r^2} [(\partial_r \omega)^2 - (\partial_t \omega)^2] - \frac{1}{r} \partial_r f = \\ & \frac{8\pi G}{r^2 \Phi_1^4} \left[(\Phi_1^4 r^2 - f^2 \Phi_2^2 W^2) ((\partial_r \Phi_1)^2 - (\partial_t \Phi_1)^2) + (\Phi_1^4 r^2 - f^2 \Phi_1^2 W^2) ((\partial_r \Phi_2)^2 - (\partial_t \Phi_2)^2) \right. \\ & \quad + f^2 \Phi_1^2 (\Phi_1^2 + \Phi_2^2) ((\partial_t W)^2 - (\partial_r W)^2) + \Phi_1^2 ((\partial_t \omega)^2 - (\partial_r \omega)^2) \\ & \quad + 2f^2 \Phi_1^3 (\Phi_1^2 + \Phi_2^2) (\partial_t \omega \partial_t W - \partial_r \omega \partial_r W) + 2f^2 \Phi_2^3 \Phi_1 W (\partial_r W \partial_r \Phi_1 - \partial_t W \partial_t \Phi_1) \\ & \quad + 2f^2 \Phi_1^3 \Phi_2 W (\partial_t \Phi_2 \partial_t \omega - \partial_r \Phi_2 \partial_r \omega) + 2f^2 \Phi_2^3 \Phi_2 W (\partial_t W \partial_t \Phi_2 - \partial_r W \partial_r \Phi_2) \\ & \quad + 2f^2 \Phi_1 \Phi_2 W^2 (\partial_r \Phi_2 \partial_r \Phi_1 - \partial_t \Phi_2 \partial_t \Phi_1) + 2f^2 \Phi_2^2 \Phi_1^2 W (\partial_r \Phi_1 \partial_r \omega - \partial_t \Phi_1 \partial_t \omega) \\ & \quad \left. + g \Phi_1^2 (\Phi_1^2 r^2 - f^2 W^2) (2A_1 (\Phi_2 \partial_r \Phi_1 - \Phi_1 \partial_r \Phi_2) + 2A_2 (\Phi_1 \partial_t \Phi_2 - \Phi_2 \partial_t \Phi_1) + g(\Phi_1^2 + \Phi_2^2)(A_1^2 - A_2^2)) \right] \quad (18) \end{aligned}$$

$$\begin{aligned} & \partial_t^2 \omega - \partial_r^2 \omega + \frac{2}{f} (\partial_t f \partial_t \omega - \partial_r f \partial_r \omega) + \frac{1}{r} \partial_r \omega = \frac{16\pi G}{f \Phi_1^2} \left[W \Phi_1 ((\partial_r \Phi_2)^2 - (\partial_t \Phi_2)^2) \right. \\ & \quad + \Phi_2 W (\partial_t \Phi_2 \partial_t \Phi_1 - \partial_r \Phi_2 \partial_r \Phi_1) + \Phi_2 \Phi_1^2 (\partial_r \Phi_2 \partial_r \omega - \partial_t \Phi_2 \partial_t \omega) + \Phi_2 \Phi_1 (\partial_r \Phi_2 \partial_r W - \partial_t \Phi_2 \partial_t W) \\ & \quad + \Phi_1^2 (\partial_r \Phi_1 \partial_r W - \partial_t \Phi_1 \partial_t W) + \Phi_1^3 (\partial_r \Phi_1 \partial_r \omega - \partial_t \Phi_1 \partial_t \omega) \\ & \quad \left. + g^2 \Phi_1 W (\Phi_1^2 + \Phi_2^2)(A_1^2 - A_2^2) + 2g \Phi_1^2 W (A_2 \partial_t \Phi_2 - A_1 \partial_r \Phi_2) + 2g \Phi_1 \Phi_2 W (A_1 \partial_r \Phi_1 - A_2 \partial_t \Phi_1) \right], \quad (19) \end{aligned}$$

and from the YM equations Eq. (4)

$$\begin{aligned} & \partial_t^2 \Phi_1 - \partial_r^2 \Phi_1 - \frac{1}{r} \partial_r \Phi_1 + \frac{f^2 \Phi_1}{r^2} ((\partial_r \omega)^2 - (\partial_t \omega)^2) + \frac{1}{f} (\partial_r f \partial_r \Phi_1 - \partial_t f \partial_t \Phi_1) \\ & \quad + \frac{f^2}{r^2} (\partial_r \omega \partial_r W - \partial_t \omega \partial_t W) + \frac{gf^2 \Phi_2 W}{\Phi_1 r^2} (A_1 \partial_r \omega - A_2 \partial_t \omega) \\ & \quad + \frac{g \Phi_2}{f} (A_1 \partial_r f - A_2 \partial_t f) + 2g (A_2 \partial_t \Phi_2 - A_1 \partial_r \Phi_2) - \frac{g A_1 \Phi_2}{r} + g^2 \Phi_1 (A_1^2 - A_2^2) = 0 \quad (20) \end{aligned}$$

$$\begin{aligned} & \partial_t^2 \Phi_2 - \partial_r^2 \Phi_2 - \frac{1}{r} \partial_r \Phi_2 + \frac{f^2 \Phi_2}{r^2} ((\partial_r \omega)^2 - (\partial_t \omega)^2) + \frac{1}{f} (\partial_r f \partial_r \Phi_2 - \partial_t f \partial_t \Phi_2) + \frac{f^2 \Phi_2}{r^2 \Phi_1} (\partial_r \omega \partial_r W - \partial_t \omega \partial_t W) \\ & \quad + \frac{f^2 \Phi_2 W}{r^2 \Phi_1^2} (\partial_t \omega \partial_t \Phi_1 - \partial_r \omega \partial_r \Phi_1) + \frac{f^2 W}{r^2 \Phi_1} (\partial_r \omega \partial_r \Phi_2 - \partial_t \omega \partial_t \Phi_2) + 2g (A_1 \partial_r \Phi_1 - A_2 \partial_t \Phi_1) \\ & \quad + \frac{g \Phi_1}{f} (A_2 \partial_t f - A_1 \partial_r f) + \frac{gf^2 W}{r^2} (A_2 \partial_t \omega - A_1 \partial_r \omega) + g^2 \Phi_2 (A_1^2 - A_2^2) + \frac{g A_1 \Phi_1}{r} = 0 \quad (21) \end{aligned}$$

$$\begin{aligned} & \partial_t^2 W_1 - \partial_r^2 W_1 + \frac{1}{r} \partial_r W + \frac{1}{f} (\partial_t f \partial_t W - \partial_r f \partial_r W) + \frac{\Phi_1}{f} (\partial_r f \partial_r \omega - \partial_t f \partial_t \omega) \\ & \quad + \partial_t \omega \partial_t \Phi_1 - \partial_r \omega \partial_r \Phi_1 + \frac{2g \Phi_2}{\Phi_1} (A_2 \partial_t W - A_1 \partial_r W) + g \Phi_2 (A_2 \partial_t \omega - A_1 \partial_r \omega) \\ & \quad + \frac{2g W \Phi_2}{\Phi_1^2} (A_1 \partial_r \Phi_1 - A_2 \partial_t \Phi_1) + \frac{2g W}{\Phi_1} (A_2 \partial_t \Phi_2 - A_1 \partial_r \Phi_2) + \frac{g W \Phi_2}{f \Phi_1} (A_2 \partial_t f - A_1 \partial_r f) \\ & \quad + g^2 W (A_1^2 - A_2^2) + \frac{g A_1 \Phi_2 W}{r \Phi_1} - \frac{16\pi G}{f} \left[W ((\partial_t \Phi_2)^2 - (\partial_r \Phi_2)^2) \right. \\ & \quad + \Phi_1 (\partial_t \Phi_1 \partial_t W - \partial_r \Phi_1 \partial_r W) + \Phi_1^2 (\partial_t \omega \partial_t \Phi_1 - \partial_r \omega \partial_r \Phi_1) \\ & \quad + \Phi_2 (\partial_t \Phi_2 \partial_t W - \partial_r \Phi_2 \partial_r W) + \frac{W \Phi_2}{\Phi_1} (\partial_r \Phi_2 \partial_r \Phi_1 - \partial_t \Phi_2 \partial_t \Phi_1) + \Phi_1 \Phi_2 (\partial_t \Phi_2 \partial_t \omega - \partial_r \Phi_2 \partial_r \omega) \\ & \quad \left. + 2g W \Phi_1 (A_1 \partial_r \Phi_2 - A_2 \partial_t \Phi_2) + 2g W \Phi_2 (A_2 \partial_t \Phi_1 - A_1 \partial_r \Phi_1) + g^2 \Phi_1^2 W (\Phi_1^2 + \Phi_2^2)(A_2^2 - A_1^2) \right] = 0. \quad (22) \end{aligned}$$

III. NUMERICAL SOLUTION

In the vacuum situation, the behavior of the cylindrical wave solution is well established [39,40]. One can have reflection of incoming into outgoing waves with the same polarization. Further, there is a non-linear effect describing the interaction between the two different polarization + and × modes. If an outgoing cylindrical wave is linear polarized, its polarization vector rotates as it propagates. This effect is often called 'Faraday' rotation and is given by

$$\tan \theta_A = \frac{A_x}{A_+}, \quad \tan \theta_B = \frac{B_x}{B_+} \quad (23)$$

with

$$A_{+,x} = \frac{1}{f}(\partial_t f \pm \partial_r f), \quad B_{+,x} = \frac{f}{r}(\partial_t \omega \pm \partial_r \omega) \quad (24)$$

A wave emitted from the axis of symmetry behaves like a shock front and is given by

$$f = w(a^2 + 1)/(a^2 + w^2), \quad \omega = ra(1 - w^2)/w(a^2 + 1), \quad \Omega^2 f = b\sqrt{r}/\sqrt{t^2 - r^2}, \quad (25)$$

with $w = (t - \sqrt{t^2 - r^2})/r$ and a and b arbitrary constants. One can generate a two-soliton solution and it is found that near the axis $\log(\Omega^2 f)$, which also describes the 'C-energy', does not approach zero. So some matter field must act as a possible source of the two-soliton field. This curious conclusion can also be formulated in context of the cosmic string with mass unit length $m = p/(2p - 2)$, with p a constant originating from the static solution $f = r^{2p}$, $\Omega = ar^{p^2-p}$, which on his turn is related to the angle deficit of the string, given by

$$\delta\phi = 2\pi(1 - \frac{1}{\sqrt{\Omega^2 f}}) \quad (26)$$

When a 'Rosen'-pulse $\ln f = \int_0^{t-r} f(\beta) d\beta / \sqrt{(t-\beta)^2 - r^2}$ is emitted, $\log(\Omega^2 f)$ changes by an amount $-k^2$, where k depends solely on $f(\beta)$. So after the pulse, one has a different string, which let Marder [41] to conclude that the mass per unit length has decreased by an amount $\sim k^2$. Because $\log(\Omega^2 f)$ is related to the 'C'-energy, this confirms that the waves carry away energy and are evidently generated by a source on the axis. So a continuous change in the parameter p may produce suddenly and inexplicable change in physical meaning.

Further, it was emphasized in the vacuum case that there is a reflection of the outgoing wave into an ingoing wave and visa versa.

From the Einstein-Maxwell case [21,23] it is known that there exist stable open cosmic strings coupled with gravity and electromagnetism. When gravitational and electromagnetic waves are generated, the angle deficit is increased asymptotically and near the axis independent of the coupling of the Maxwell field to gravity. Further, it was emphasized that the electromagnetic waves do contribute to the rotation of the spacetime by effecting the 'dragging' of the Killing field $\frac{\partial}{\partial z}$, but not to the angle deficit.

In our EYM case we solved Eq.(17)-(22) for the initial values

$$\Omega(0, r) = C_1 r^{p^2-p}, \quad f(0, r) = r^{2p}, \quad \Phi_1(0, r) = 1 - e^{-C_2 r}, \quad \Phi_2(0, r) = e^{-C_3 r} - 1 \quad (27)$$

For W_1 we will take as initial value a node number 1 or node number 2 function, i.e., $W_1(0, r) = -\tanh(r - 1)$, or $W_1(0, r) = 1 - e^{-C_4 r^2}$. Here are C_i some constants.

We used a 100x100 grid and applied the method of lines with cubic Hermite polynomials. The roundoff error remained below 0.001. In figure 1 we took $\omega(0, r) = \frac{3.5r}{r^2+1}$ and for W_1 the node number 2 function. We plotted respectively: Ω , f , ω , Φ_1 , W_1 , Φ_2 , the angle deficit, the gravitational puls wave $\Omega^2 f$, $g_{\varphi\varphi}$, and W_2 . In figure 2 we took a different initial Ω (p=0.1 in stead of 0.25). We obtain a solitonlike solution. The behavior depends on the initial mass per unit length of the string, just as in the vacuum case. In figure 3 we took the node number 1 function for W_1 . We observe that the angle deficit changes significantly as well as the gravitational outgoing pulse. So the Yang-Mills waves has an impact on the gravitational waves. In figure 4 we took p=0.1 in stead of p=0.25. We see that the behavior of the solution changes significantly: the angle deficit close to the r-axis increases, while the gravitational puls 'hangs' close to the r-axes.

In figure 5 we plotted a long-time run for a different initial ω : an outgoing wave is reflected into an ingoing wave.

An important question is if the solution admits gauge-charges. The gauge-charges Q_E, Q_M (electric and magnetic) are given by

$$Q_E = \frac{1}{4\pi} \oint_S |^*F|, \quad Q_M = \frac{1}{4\pi} \oint_S |F|, \quad (28)$$

with F the YM field strength, $^*F$ the dual tensor and the integration over a two-sphere at spatial infinity. In our situation, we obtain for the charges

$$Q_E = \int r \left\{ \left[\partial_t(W_1 + \omega\Phi_1) + gA_2(W_2 - 1 + \omega\Phi_2) + A_2 \right] \tau_r + \left[\partial_t(W_2 + \omega\Phi_2) - gA_2(W_1 + \omega\Phi_1) \right] \tau_z \right\} dr \quad (29)$$

$$Q_M = \int \frac{r}{\Omega^2} \left\{ \left[\frac{1}{f} (\partial_r \Phi_1 + gA_1 \Phi_2 - \frac{f\omega}{r^2} (\Phi_1 \partial_r \omega + \partial_r W_1 + A_1 + gA_1(W_2 - 1))) \tau_r \right] + \left[\frac{1}{f} (\partial_r \Phi_2 - gA_1 \Phi_1) - \frac{f\omega}{r^2} (\partial_r W_2 + \Phi_2 \partial_r \omega - gA_1 W_1) \right] \tau_z \right\} dr, \quad (30)$$

where we used Eq. (15). So in general the solution will possess gauge charges.

IV. AN APPROXIMATE WAVE SOLUTION TO FIRST ORDER

In the appendix we presented the multiple-scale method for the EYM system. On the metric Eq. (7) we have for the tetrad component: $l_\mu = (-1, 1, 0, 0)$ and $l^\mu = \frac{1}{\Omega^2}(1, 1, 0, 0)$, which implies that the divergence of the null congruence l^μ becomes

$$\nabla_\mu l^\mu = \frac{1}{\Omega^2 r}, \quad (31)$$

as it should be. From Eq. (A10) we obtain $B_0^a = -B_1^a$, or, in the notation of Eq. (14), $\dot{A}_1^{(1)} = -\dot{A}_2^{(1)}$. From Eq. (A8) we obtain

$$\dot{h}_{01} = -\frac{1}{2}(\dot{h}_{00} + \dot{h}_{11}), \quad \dot{h}_{12} = -\dot{h}_{02}, \quad \dot{h}_{03} = -\dot{h}_{13}, \quad \dot{h}_{23} = -\frac{\bar{g}^{22}\dot{h}_{22} + \bar{g}^{33}\dot{h}_{33}}{2\bar{g}^{23}}. \quad (32)$$

First of all we have just as Eq. (15),

$$\partial_t \bar{A}_1 = \partial_r \bar{A}_2, \quad \bar{W}_2 = \frac{\bar{\Phi}_2 \bar{W}_1}{\bar{\Phi}_1} + 1 - \frac{1}{g} \quad (33)$$

For simplicity we will take $\Phi_2 = 0$ and try to find an approximate wave solution to second order. From the propagation equations Eq. (A14) and (A18) we obtain

$$\bar{\Phi}_1 \partial_\varphi \dot{\omega}^{(1)} + \partial_\varphi \dot{W}_1^{(1)} = 0, \quad (34)$$

$$\partial_\varphi \dot{W}_2^{(1)} = 0. \quad (35)$$

So we will take $\dot{\omega}^{(1)}$ and $\dot{W}_2^{(1)}$ independent of φ . Further,

$$\partial_t \dot{A}_1^{(1)} + \partial_r \dot{A}_1^{(1)} = \ddot{A}_1^{(2)} + \ddot{A}_2^{(2)} - \frac{g\bar{f}\bar{\Omega}^2}{r^2} \bar{W}_1 \dot{W}_2^{(1)}, \quad (36)$$

$$\begin{aligned} \partial_t \dot{\Phi}_1^{(1)} + \partial_r \dot{\Phi}_1^{(1)} &= -\frac{\dot{\Phi}_1^{(1)}}{2r} + \frac{\dot{f}^{(1)}}{2f} (\partial_t \bar{\Phi}_1 + \partial_r \bar{\Phi}_1) + \frac{\dot{\Phi}_1^{(1)}}{2f} (\partial_t \bar{f} + \partial_r \bar{f}) \\ &+ \frac{\bar{f}^2}{r^2} (\dot{\omega}^{(1)} \bar{\Phi}_1 + \frac{1}{2} \dot{W}_1^{(1)}) (\partial_t \bar{\omega} + \partial_r \bar{\omega}) + \frac{\bar{f}^2 \dot{\omega}^{(1)}}{2r^2} (\partial_t \bar{W}_1 + \partial_r \bar{W}_1), \end{aligned} \quad (37)$$

$$\partial_t \dot{W}_2^{(1)} + \partial_r \dot{W}_2^{(1)} = -\frac{1}{2f} \dot{W}_2^{(1)} (\partial_t \bar{f} + \partial_r \bar{f}) + \frac{1}{2r} \dot{W}_2^{(1)} + g(\bar{A}_1 + \bar{A}_2) \left(\dot{W}_1^{(1)} + \frac{\bar{\Phi}_1 \dot{\omega}^{(1)}}{2} + \frac{\dot{f}^{(1)} \bar{W}_1}{2f} \right) \quad (38)$$

$$\begin{aligned}\partial_t \dot{W}_1^{(1)} + \partial_r \dot{W}_1^{(1)} &= \frac{1}{2r} \dot{W}_1^{(1)} - \frac{1}{2\bar{f}} (\dot{W}_1^{(1)} + \bar{\Phi}_1 \dot{\omega}^{(1)}) (\partial_t \bar{f} + \partial_r \bar{f}) - \bar{\Phi}_1 (\partial_t \dot{\omega}^{(1)} + \partial_r \dot{\omega}^{(1)}) - \frac{\dot{f}^{(1)}}{2\bar{f}} (\partial_r \bar{W}_1 + \partial_t \bar{W}_1) \\ &\quad - \frac{1}{2} (\partial_t \bar{\omega} + \partial_r \bar{\omega}) (\dot{\Phi}_1^{(1)} + \frac{\bar{\Phi}_1 \dot{f}^{(1)}}{\bar{f}}) - \frac{\dot{\omega}^{(1)}}{2} (\partial_t \bar{\Phi}_1 + \partial_r \bar{\Phi}_1) + \frac{\bar{\Phi}_1 \dot{\omega}^{(1)}}{2r} - g \dot{W}_2^{(1)} (\bar{A}_1 + \bar{A}_2)\end{aligned}\quad (39)$$

$$\begin{aligned}\partial_t \dot{\Omega}^{(1)} + \partial_r \dot{\Omega}^{(1)} &= -\frac{\bar{\Omega} \dot{f}^{(1)}}{4\bar{f}^2} (\partial_t \bar{f} + \partial_r \bar{f}) + \frac{\dot{\Omega}^{(1)}}{\bar{\Omega}} (\partial_t \bar{\Omega} + \partial_r \bar{\Omega}) + \frac{\bar{\Omega} \dot{f}^{(1)}}{4r\bar{f}} + \frac{1}{4\bar{\Omega}} (\ddot{k}_{00} + \ddot{k}_{11} + 2\ddot{k}_{01}) \\ &\quad - \frac{8\pi G \bar{\Omega} \bar{f}}{r^2} \left[\frac{\dot{\Phi}_1^{(1)}}{\bar{f}^2} (\partial_t \bar{\Phi}_1 + \partial_r \bar{\Phi}_1) + (\dot{\omega}^{(1)} \bar{\Phi}_1 + \dot{W}_1^{(1)}) (\partial_t \bar{W}_1 + \partial_r \bar{W}_1 + \bar{\Phi}_1 (\partial_t \bar{\omega} + \partial_r \bar{\omega})) - g \dot{W}_2^{(1)} \bar{W}_1 (\bar{A}_1 + \bar{A}_2) \right]\end{aligned}\quad (40)$$

$$\begin{aligned}\partial_t \dot{f}^{(1)} + \partial_r \dot{f}^{(1)} &= \frac{\dot{f}^{(1)}}{\bar{f}} (\partial_t \bar{f} + \partial_r \bar{f}) - \frac{\dot{f}^{(1)}}{2r} + \frac{\bar{f}^3 (\partial_t \bar{\omega} + \partial_r \bar{\omega}) \dot{\omega}^{(1)}}{r^2} - \frac{8\pi G \bar{f}^2}{r^2} \left((\dot{\omega}^{(1)} \bar{\Phi}_1 + \dot{W}_1^{(1)}) (\partial_t \bar{W}_1 + \partial_r \bar{W}_1) \right. \\ &\quad \left. + (\bar{\Phi}_1^2 \dot{\omega}^{(1)} + \bar{\Phi}_1 \dot{W}_1^{(1)}) (\partial_t \bar{\omega} + \partial_r \bar{\omega}) + \frac{3r^2 \dot{\Phi}_1^{(1)}}{\bar{f}^2} (\partial_t \bar{\Phi}_1 + \partial_r \bar{\Phi}_1) \right),\end{aligned}\quad (41)$$

and

$$\begin{aligned}\partial_t \dot{\omega}^{(1)} + \partial_r \dot{\omega}^{(1)} &= \frac{1}{2r} \dot{\omega}^{(1)} - \frac{\dot{f}^{(1)}}{\bar{f}} (\partial_t \bar{\omega} + \partial_r \bar{\omega}) - \frac{\dot{\omega}^{(1)}}{\bar{f}} (\partial_t \bar{f} + \partial_r \bar{f}) - \frac{8\pi G}{\bar{f}} \left(\bar{\Phi}_1 \dot{\Phi}_1^{(1)} (\partial_t \bar{\omega} + \partial_r \bar{\omega}) \right. \\ &\quad \left. + (\bar{\Phi}_1 \dot{\omega}^{(1)} + \dot{W}_1^{(1)}) (\partial_t \bar{\Phi}_1 + \partial_r \bar{\Phi}_1) + \dot{\Phi}_1^{(1)} (\partial_t \bar{W}_1 + \partial_r \bar{W}_1) - g \bar{\Phi}_1 \dot{W}_2^{(1)} (\bar{A}_1 + \bar{A}_2) \right)\end{aligned}\quad (42)$$

These perturbation equations are linear differential equations of first order in $\dot{h}_{\mu\nu}$ and $\dot{B}_\mu^a$ and contain in general the term $\ddot{k}_{\mu\nu}$ and $\ddot{C}_\mu^a$. In our special case, we see appear in Eq. (38) a second order term $\ddot{k}_{\mu\nu}$ and in Eq. (34) a term $\ddot{A}_i^{(2)}$. The remaining perturbation equations will also contain second order term like $\ddot{W}_i^{(2)}$.

For the background variables $\bar{\Phi}_1, \bar{W}_1$ and $\bar{W}_2$ we use Eq.(A13). The resulting equations are similar to the equations found in section 2 for $\Phi_2 = 0$. From Eq. (A17) we obtain equations for the background variables $\bar{\Omega}, \bar{f}$, and $\bar{\omega}$ with back-reaction terms, for example,

$$\begin{aligned}\partial_t^2 \bar{\Omega} - \partial_r^2 \bar{\Omega} &= -\frac{1}{r} \partial_r \bar{\Omega} + \frac{1}{\bar{\Omega}} \left((\partial_t \bar{\Omega})^2 - (\partial_r \bar{\Omega})^2 \right) + \frac{\bar{\Omega} (\partial_r \bar{f})^2}{2\bar{f}^2} - \frac{\bar{\Omega} \partial_r \bar{f}}{\bar{f} r} + \frac{\bar{f}^2 \bar{\Omega} (\partial_r \bar{\omega})^2}{2r^2} \\ &\quad + \frac{\bar{\Omega}}{2\tau \bar{f}^2 r^2} \int [\bar{f}^4 (\dot{\omega}^{(1)})^2 + r^2 (\dot{f}^{(1)})^2] d\xi + \frac{8\pi G \bar{\Omega}}{\bar{f} r^2 \tau} \int [\bar{f}^2 (\dot{W}_1^{(1)} + \bar{\Phi}_1 \dot{\omega}^{(1)})^2 + r^2 (\dot{\Phi}_1^{(1)})^2 + \bar{f}^2 (\dot{W}_2^{(1)})^2] d\xi \\ &\quad + \frac{4\pi G \bar{\Omega} \bar{f}}{r^2} \left[(\partial_t \bar{W}_1 + \bar{\Phi}_1 \partial_t \bar{\omega})^2 + (\partial_r \bar{W}_1 + \bar{\Phi}_1 \partial_r \bar{\omega})^2 + \frac{r^2}{\bar{f}^2} (\partial_r \bar{\Phi}_1^2 + \partial_t \bar{\Phi}_1^2) \right].\end{aligned}\quad (43)$$

So we see that a first order HF gravitational and Yang-Mills wave emitted from the string, will change the metric background component $\bar{\Omega}$, just as in the vacuum situation for pure gravitational waves. Moreover, there will be a distortion of the shape of the wave during its propagation, which does not occur in the pure Einstein case. In general, this is due to the fact that second order terms appear in the perturbation equations of $\dot{h}_{\mu\nu}$ and $\dot{B}_\mu^a$. Now let us consider, for example, the equation for $\dot{W}_2^{(1)}$, Eq. (36). For $A_1 = -A_2$, we can solve $\dot{W}_2^{(1)}$ for $\bar{f} = e^{2p}$. For example for $p = 1/4$ we have

$$\dot{W}_2^{(1)} = \sqrt{r} B(t-r) e^{-2/3r^{1.5}}, \quad (44)$$

which approaches to zero for large r . Here is $B(t-r)$ an arbitrary function of $(t-r)$. Together with Eq.(31) we then have to first order

$$\dot{W}_2^{(1)} = 1 - 1/g + \sqrt{r} B(t-r) e^{-2/3r^{1.5}} + \dots, \quad (45)$$

which is bounded for large r . One can proceed in the same way for the other first order perturbations by imposing some trial solutions for the background variables [10]

V. CONCLUSIONS

Non-Abelian gravity-coupled solitons and black holes could have played an important role in gauge theories of elementary particles in the early stage of the universe. Besides the well-known spherically symmetric solutions, axially symmetric solutions are of interest due to the fact that they admit cosmic strings, which could have played the role of 'seeds' for the large-scale structure of clusters in the universe. It is quite evident that the solution found here is a radiating one and there is an interaction of the solitonlike gravitational waves and the Yang-Mills waves with the rotating string. This can be seen from figure 5 where the angle deficit and Ω is significantly changed by the pulses. This can also be seen from Eq. (41), where the term on the right-hand side represents the back-reaction of the waves on Ω . The wave-like solutions found in this model is significantly different from the electrovac solution found in the Einstein-Maxwell model. The intervening electro-magnetic waves did not contribute to the angle deficit asymptotically, while in our model the magnetic components W_i has an impact on the angle deficit for large r (compare figures 1 and 3). In particular the number of times the W_i crosses the r -axis influences the gravitational wave puls. Further, for a different mass per unit length of the string, a different behavior of the solution is obtained (compare figures 3 and 4).

In order to conclude if the solution is topological stable, we must solve the perturbation equations to higher order. Moreover, a combination of the results of a throughout numerical investigation with the multiple-scale method pushed to higher orders, will lead to further understanding of the rich structure of the EYM system on a rotating space time. Specially the question if a electrically and magnetically charged solution ('dyon') of the rotating cosmic string exists. Further, bursts of high-frequency gravitational waves might be detectable by the planned gravitational wave detectors LIGO/VIRGO and LISA, even for strings at GUT scale [42].

The issues mentioned above are currently under study by the author.

APPENDIX A: THE EYM EQUATIONS IN THE MULTIPLE SCALE FORMULATION

Let us consider a manifold $\mathcal{M}$ with two different scales, i.e., a mapping from $\mathcal{M} \times \mathcal{R}$ into the space of metrics on $\mathcal{M}$:

$$(\mathbf{x}, \xi) \implies g(\mathbf{x}, \xi), \quad \mathbf{x} \in \mathcal{M}, \xi \in \mathcal{R}. \quad (\text{A1})$$

We set $\xi \equiv \omega \Pi(\mathbf{x})$, with Π a phase function on $\mathcal{M}$ of dimension of *length* and ω a large parameter of dimension $(\text{length})^{-1}$.

The multiple-scale method assumes that the metric $g_{\mu\nu}$ and the YM-potentials A_μ^a can be written as

$$g_{\mu\nu} = \bar{g}_{\mu\nu} + \frac{1}{\omega} h_{\mu\nu}(x^\sigma; \xi) + \frac{1}{\omega^2} k_{\mu\nu}(x^\sigma; \xi) + \dots, \quad (\text{A2})$$

$$A_\mu^a = \bar{A}_\mu^a + \frac{1}{\omega} B_\mu^a(x^\sigma; \xi) + \frac{1}{\omega^2} C_\mu^a(x^\sigma; \xi) + \dots, \quad (\text{A3})$$

where $\xi \equiv \omega \Pi(x^\sigma)$ and Π a phase function. The parameter ω measures the ratio of the fast scale to the slow one. The rapid variation only occur in the direction of the vector $l_\sigma \equiv \frac{\partial \Pi}{\partial x^\sigma}$. For a function $\Psi(x^\sigma; \xi)$ one has

$$\frac{\partial \Psi}{\partial x^\sigma} = \partial_\sigma \Psi + \omega l_\sigma \dot{\Psi}, \quad (\text{A4})$$

where $\partial_\sigma \Psi \equiv \frac{\partial \Psi}{\partial x^\sigma}|_{\xi \text{ fixed}}$ and $\dot{\Psi} \equiv \frac{\partial \Psi}{\partial \xi}|_{x^\sigma \text{ fixed}}$. We consider here the case where the magnitude of the perturbation $h_{\mu\nu}$ of the metric with respect to the background $\bar{g}_{\mu\nu}$ is of order ω^{-1} (for different possibilities, such as for example with the leading term in the metric expansion of order ω^{-2} , see [13]). Further, we will consider here the hypersurfaces $\Pi(x^\sigma) = \text{cst}$ as wave fronts for the background metric $\bar{g}$, so (Eikonal equation)

$$l_\alpha l_\beta \bar{g}^{\alpha\beta} = 0. \quad (\text{A5})$$

Substituting the expansions of the field variables into the equations and collecting terms of equal orders of ω , one obtains propagation equations for $\dot{B}_\mu^a, \dot{C}_\mu^a, \dot{h}_{\mu\nu}$ and $\dot{k}_{\mu\nu}$ and 'back-reaction' equations for $\bar{g}_{\mu\nu}$ and $\bar{A}_\mu^a$. It will be clear from the propagation equation that there will be a coupling between the high-frequency gravitational field and the high-frequency behavior of A_μ^a when the singularity will be approached.

First we substitute the expansions Eq. (A2) and (A3) into the YM-equation Eq. (4). We obtain for the order ω equation

$$\bar{g}^{\mu\alpha} [l_\mu l_\alpha \ddot{B}_\nu^a - l_\nu l_\alpha \ddot{B}_\mu^a] = 0. \quad (\text{A6})$$

For the order ω^0 -equation we obtain

$$\begin{aligned} \bar{g}^{\mu\alpha} \Big[& \bar{\nabla}_\alpha \bar{\mathcal{F}}_{\mu\nu}^a + \Upsilon_{\alpha\mu}^\lambda \bar{\mathcal{F}}_{\nu\lambda}^a - \Upsilon_{\alpha\nu}^\lambda \bar{\mathcal{F}}_{\mu\lambda}^a + \dot{B}_\nu^a (l_{\mu,\alpha} - (\bar{\Gamma}_{\alpha\mu}^\lambda + \Upsilon_{\alpha\mu}^\lambda) l_\lambda) - \dot{B}_\mu^a (l_{\nu,\alpha} - (\bar{\Gamma}_{\alpha\nu}^\lambda + \Upsilon_{\alpha\nu}^\lambda) l_\lambda) \\ & + l_\mu (\dot{B}_{\nu,\alpha} - (\bar{\Gamma}_{\alpha\nu}^\lambda + \Upsilon_{\alpha\nu}^\lambda) \dot{B}_\lambda^a) - l_\nu (\dot{B}_{\mu,\alpha} - (\bar{\Gamma}_{\alpha\mu}^\lambda + \Upsilon_{\alpha\mu}^\lambda) \dot{B}_\lambda^a) + l_\alpha (\dot{B}_{\nu,\mu}^a - \dot{B}_{\mu,\nu}^a) \\ & + l_\alpha (l_\mu \ddot{C}_\nu^a - l_\nu \ddot{C}_\mu^a) + g\epsilon^{abc} (l_\alpha (\bar{A}_\mu^b \dot{B}_\nu^c + \bar{A}_\nu^c \dot{B}_\mu^b) + \bar{A}_\alpha^b (\bar{\mathcal{F}}_{\mu\nu}^c + l_\mu \dot{B}_\nu^c - l_\nu \dot{B}_\mu^c)) \Big] \\ & - h^{\mu\alpha} l_\alpha (l_\mu \ddot{B}_\nu^a - l_\nu \ddot{B}_\mu^a) = 0, \end{aligned} \quad (\text{A7})$$

with $\Upsilon_{\mu\nu}^\lambda \equiv \frac{1}{2} \bar{g}^{\sigma\lambda} (l_\mu \dot{h}_{\nu\sigma} - l_\nu \dot{h}_{\mu\sigma} - l_\sigma \dot{h}_{\mu\nu})$.

Substituting the expansions into the Einstein equations Eq. (3), we obtain for the order ω equation

$$R_{\mu\nu}^{(-1)} = l_\nu \dot{\Upsilon}_{\mu\sigma}^\sigma - l_\sigma \dot{\Upsilon}_{\mu\nu}^\sigma = 0. \quad (\text{A8})$$

For the ω^0 -equation we obtain

$$\begin{aligned} \bar{R}_{\mu\nu} + R_{\mu\nu}^{(0)} = & -8\pi G \left\{ \bar{g}^{\lambda\beta} (\bar{\mathcal{F}}_{\mu\lambda}^a + l_\mu \dot{B}_\lambda^a - l_\lambda \dot{B}_\mu^a) (\bar{\mathcal{F}}_{\nu\beta}^a + l_\nu \dot{B}_\beta^a - l_\beta \dot{B}_\nu^a) \right. \\ & \left. + \frac{1}{4} \bar{g}_{\mu\nu} \bar{g}^{\sigma\alpha} \bar{g}^{\tau\beta} (\bar{\mathcal{F}}_{\alpha\beta}^a + l_\alpha \dot{B}_\beta^a - l_\beta \dot{B}_\alpha^a) (\bar{\mathcal{F}}_{\sigma\tau}^a + l_\sigma \dot{B}_\tau^a - l_\tau \dot{B}_\sigma^a) \right\}, \end{aligned} \quad (\text{A9})$$

with $\bar{R}_{\mu\nu}$ the background Ricci tensor and $R_{\mu\nu}^{(0)}$ an expression in $\ddot{k}, h\ddot{h}, \dot{h}^2$ and $\dot{h}$ (see [11]). We can simplify the equations considerable. If we use Eq. (A5), then Eq. (A6) becomes

$$l^\mu \ddot{B}_\mu^a = 0, \quad (\text{A10})$$

so for periodic B_μ^a we have

$$l^\mu B_\mu^a = 0. \quad (\text{A11})$$

Using Eq. (A8) and Eq. (A10) we obtain from Eq. (A7)

$$\begin{aligned} \bar{g}^{\mu\alpha} \Big[& \bar{\nabla}_\alpha \bar{\mathcal{F}}_{\mu\nu}^a - \Upsilon_{\alpha\nu}^\lambda \bar{\mathcal{F}}_{\mu\lambda}^a + \dot{B}_\nu^a \bar{\nabla}_\alpha l_\mu - \dot{B}_\mu^a \bar{\nabla}_\alpha l_\nu + l_\mu \bar{\nabla}_\alpha \dot{B}_\nu^a - l_\nu \bar{\nabla}_\alpha \dot{B}_\mu^a + l_\alpha (\dot{B}_{\nu,\mu}^a - \dot{B}_{\mu,\nu}^a) \Big] \\ & + l_\nu l_\alpha h^{\mu\alpha} \ddot{B}_\mu^a - l_\nu l^\mu \ddot{C}_\mu^a + g\epsilon^{abc} \bar{g}^{\mu\alpha} [\bar{A}_\alpha^b \bar{\mathcal{F}}_{\mu\nu}^c + 2l_\alpha \bar{A}_\mu^b \dot{B}_\nu^c - l_\nu \bar{A}_\alpha^b \dot{B}_\mu^c] = 0, \end{aligned} \quad (\text{A12})$$

with $\bar{\nabla}$ the covariant derivative with respect to the background metric $\bar{g}_{\mu\nu}$. Integrating this equation with respect to ξ yields

$$\bar{\mathcal{D}}^\mu \bar{\mathcal{F}}_{\mu\nu}^a = 0. \quad (\text{A13})$$

Substituting back this equation into Eq. (A12) we obtain the propagation equation for the YM-field

$$\begin{aligned} \bar{\nabla}^\mu (l_\mu \dot{B}_\nu^a - l_\nu \dot{B}_\mu^a) + l^\mu (\bar{\nabla}_\mu \dot{B}_\nu^a - \bar{\nabla}_\nu \dot{B}_\mu^a) - \bar{g}^{\mu\alpha} \Upsilon_{\alpha\nu}^\lambda \bar{\mathcal{F}}_{\mu\lambda}^a + l^\alpha l_\nu h_\alpha^\lambda \ddot{B}_\lambda^a \\ - l_\nu l^\mu \ddot{C}_\mu^a + g\epsilon^{abc} \bar{g}^{\mu\alpha} (2l_\alpha \bar{A}_\mu^b \dot{B}_\nu^c - l_\nu \bar{A}_\alpha^b \dot{B}_\mu^c) = 0. \end{aligned} \quad (\text{A14})$$

Multiplying this propagation equation with $\dot{B}^{\nu a}$, we obtain the 'conservation'-equation

$$\bar{\nabla}_\alpha (l^\alpha \dot{B}_\nu^a \dot{B}^{\nu a}) = l^\mu \dot{h}_\nu^\lambda \dot{B}^{\nu a} \bar{\mathcal{F}}_{\mu\lambda}^a - 2g\epsilon^{abc} l^\mu \dot{B}_\nu^c \dot{B}^{\nu a} \bar{A}_\mu^b, \quad (\text{A15})$$

so $\dot{B}_\nu^a \dot{B}^{\nu a}$ is not conserved, unless the right-hand side is zero. Multiplying Eq. (A14) with l^ν , we obtain

$$l^\nu (l^\lambda \dot{h}_\nu^\mu - l^\mu \dot{h}_\nu^\lambda) \bar{\mathcal{F}}_{\mu\lambda}^a = 0. \quad (\text{A16})$$

For the choice $l^\nu h_{\nu\mu} = 0$ and hence from Eq. (A8), $h = 0$, the equations simplify again. Integrating Eq. (A9) with respect to ξ , we obtain

$$\bar{R}_{\mu\nu} = -\frac{1}{4\tau} l_\mu l_\nu \int (\dot{h}_\rho^\sigma \dot{h}_\sigma^\rho - \frac{1}{2} \dot{h}^2) d\xi - \frac{8\pi G}{\tau} l_\mu l_\nu \int \dot{B}^{\alpha a} \dot{B}_\alpha^a d\xi - 8\pi G \bar{T}_{\mu\nu}, \quad (\text{A17})$$

with τ the period of high-frequency components. In the right-hand side we have the back-reaction term due to the high-frequency gravitational perturbation, the high-frequency YM-perturbations and the background energy-momentum term $\bar{T}$. Substitution back Eq. (A17) into Eq. (A9), we obtain the perturbation equations for $h_{\mu\nu}$

$$\begin{aligned} R_{\mu\nu}^{(0)} = & l_\mu l_\nu \left[\frac{1}{4\tau} \int (\dot{h}_\rho^\sigma \dot{h}_\sigma^\rho - \frac{1}{2} \dot{h}^2) d\xi + \frac{8\pi G}{\tau} \int \dot{B}^{\alpha a} \dot{B}_\alpha^a d\xi \right] - 8\pi G \left[(l_\mu l_\nu \dot{B}^{\lambda a} \dot{B}_\lambda^a \right. \\ & \left. + \bar{\mathcal{F}}_{\mu\lambda}^a (l_\nu \dot{B}^{\lambda a} - l^\lambda \dot{B}_\nu^a) + \bar{\mathcal{F}}_{\nu\lambda}^a (l_\mu \dot{B}^{\lambda a} - l^\lambda \dot{B}_\mu^a) + \frac{1}{2} \bar{g}_{\mu\nu} \bar{F}_{\alpha\beta} (l^\alpha \dot{B}^{\beta a} - l^\beta \dot{B}^{\alpha a}) \right] \end{aligned} \quad (\text{A18})$$

-
- [1] Volkov, M. S., and Gal'tsov, D. V. 1999, Phys. Rep., 319, 1(1999)
 - [2] Bartnik, R., and McKinnon, J. 1988, Phys. Rev. Lett., 61, 141(1988)
 - [3] Smoller, J., and Wasserman, A. 1996, J. Math. Phys., 38, 6522(1996)
 - [4] Straumann, N., and Zhou, Z. H. 1990, Phys. Lett., B237, 353(1990)
 - [5] Lavrelashvili, G., and Maison, D. 1995 Phys. Lett., 343, 214(1995)
 - [6] Straumann, N., and Zhou, Z. H. 1990, Phys. Lett., B243, 33(1990)
 - [7] Zhou, Z. H., and Straumann, N. 1991, Nucl. Phys., B360, 180(1991)
 - [8] Breitenlohner, P., Lavrelashvili, G., and Maison, D. 1997, gr-qc/9708036
 - [9] Mavromatos, N. E., and Winstanley, E. 1998, J. Math. Phys., 39, 4849(1998)
 - [10] Slagter, R. J., In *Gravitational Waves and Experimental Gravity*, Proceedings of XXXIVth Rencontres de Moriond, Les Arcs, France, 1999
 - [11] Choquet-ruhat, Y. 1969, Commun. Math. Phys., 12, 16(1969)
 - [12] Choquet-Bruhat, Y. 1977, Gen. Rel. Grav., 8, 561(1977)
 - [13] Taub, A. H., in *General Relativity and Gravitation*, ed. A. Held (Plenum press, New York, 1980)
 - [14] Slagter, R. J. 1991, Gen. Rel. Grav., 23, 991(1991)
 - [15] Choptuik, M. W. 1993, Phys. Rev. Lett., 70, 9(1993)
 - [16] Choptuik, M. W., Chmaj, T., and Bizon, P. 1996, Phys. Rev. Lett., 77, 424(1996)
 - [17] Garfinkle, D. 1997, Phys. Rev., D56, R3169(1997)
 - [18] Bizon, P. 1996, gr-qc/960606(1996)
 - [19] Stachel, J. 1966, J. Math. Phys., 7, 1321(1966)
 - [20] Piran, T., Safer, P. N., and Stark, R. F. 1985, Phys. Rev., D32, 3101(1985)
 - [21] Xanthopoulos, B. C. 1986, Phys. Lett., B178, 163(1986)
 - [22] Economou, A., and Tsoubelis, D. 1988, Phys. Rev., D38, 498(1988)
 - [23] Xanthopoulos, B. C. 1987, Phys. Rev., D35, 3713(1987)
 - [24] Kramer, D., Stephani, H., and MacCallum, M. 1980, in *Exact Solutions of Einstein's Field Equations*(Cambridge Univ. Press, 1980)
 - [25] Maeda, K., Sasaki, M., Nakamura, T., and Miyama, S. 1980, Prog. Theor. Phys., 63, 719(1980)
 - [26] Slagter, R. J. 1999, Phys. Rev., D59, 59025009(1999)
 - [27] Gal'tsov, D. V. 1998, gr-qc/9808002
 - [28] Jensen, B., and Soleng, H. H. 1992, Phys. Rev., D45, 3528(1992)
 - [29] Soleng, H. H. 1992, Gen. Rel. Grav., 24, 111(1992)
 - [30] Harari, D., and Polychronakos, A. P. 1988, Phys. Rev., D38, 3320(1988)
 - [31] Deser, S., Jackiw, R., and 't Hooft, G. 1992, Phys. Rev. Lett., 68, 267(1992)
 - [32] Letelier, P. , S. 1995, Class. Quantum Grav., 12, 471(1995)
 - [33] Mazur, P. O. 1986, Phys. Rev. Lett., 57, 929(1986)
 - [34] Bonner, W. B. 1991, Phys. Lett. A., 158, 23(1991)
 - [35] Slagter, R. J. 1995, Int. J. Mod. Phys., 1995 D4, 267(1995)
 - [36] Islam, J. N. 1985, in *Rotating Fields in General Relativity*(Cambridge Univ. Press)
 - [37] Heusler, M., and Straumann, N. 1993, Class. Quant. Grav., 10, 21(1993)
 - [38] Slagter, R. J. 1996, Phys. Rev., D54, 4873(1996)

- [39] Bonner, W. B., Griffiths, J. B., and MacCallum, M. H. H. 1994, *Gen. Rel. Grav.*, 26, 687(1994)
- [40] Tomimatsu, A. 1989, *Gen. Rel. Grav.*, 21, 613(1989)
- [41] Marder, L. 1958, *Proc. Roy. Soc. Lond.*, A408, 209(1958)
- [42] Damour, T., and Vilenkin, A. 2000, [gr-qc/0004075](#)

Figure 1. Plot of the metric- and YM components for the initial values: $\Omega = Ar^{p^2-p}(A = 3, p = 0.25)$, $f = r^{2p}$, $\omega = \frac{3.5r}{r^2+1}$, $\Phi_1 = 1 - e^{-0.8r}$, $w = 0.8 - e^{-(r-2)^2}$, $\Phi_2 = e^{-0.32r} - 1$, $G=4.5$ $g=1.25$.

Figure 2. As figure 1, with $p=0.1$, $G=0.6$

Figure 3. As figure 2, for the initial W_1 : $W_1 = -\tanh(r - 1)$

Figure 4. As figure 3, for a smaller value of the mass per unit length of the string: $p=0.09$ in stead of 0.25.

Figure 5. A long time run for $\Phi_1 = 1 - e^{-0.5r}$

